

CHAMOT

**重组人成纤维生长因子-1/酸性成纤维细胞生长因子
/Recombinant FGF-1, Human, Animal-Free**

CM090-5HP

CM090-20HP

CM090-100HP

CM090-500HP

CM090-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人成纤维生长因子-1/酸性成纤维细胞生长因子 /Recombinant FGF-1, Human, Animal-Free

产品编号	CM090-5HP	CM090-20HP	CM090-100HP	CM090-500HP	CM090-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Fibroblast Growth Factors-1 (FGF-1) is a Growth Factors of mitogenic peptides which is a 15 kDa protein containing 139 amino acid residues. Fibroblast Growth Factors-1 is secreted by the macrophage which induce endothelial cell proliferation and angiogenesis. During wound injury, fibroblast Growth Factors-1 helps the tissue remodeling. When the fibroblast Growth Factors-1 binds with the FGFR on the cell membrane which is endocytosed by FGFR and induces the cell cycle.

别称

fibroblast growth factor acidic, HBGF-1, ECGF-beta, FGF1, FGF acidic, aFGF

蛋白编码

P05230

分子量

The protein has a calculated MW of 16.77 kDa. The protein migrates as 18kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <0.3 ng/mL. The specific activity of recombinant human FGF-1 is > 1 x 10⁶ IU/mg.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MFNLPPGNYKKPKLLYCSNGGHFLRILPDGTVDGTRDRSDQHIQLQLSAESVGEVYIKSTETGQ
YLAMDTDGLLYGSQTPNEECLFLERLEENHYNTYISKKHAENWFVGLKKNKGCKRGPRTHYG
QKAILFLPLPVSSD with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491