# **CHAMOT**

重组人成纤维细胞生长因子-6 /Recombinant FGF-6, Human, Animal-Free

CM095-5HP

CM095-20HP

CM095-100HP

CM095-500HP

CM095-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人成纤维细胞生长因子-6 /Recombinant FGF-6, Human, Animal-Free

产品编号	CM095-5HP	CM095-20HP	CM095-100HP	CM095-500HP	CM095-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介			
背景描述	Fibroblast Growth Factors-6 (FGF-6) is a 22.9 kDa member of the fibroblast Growth Factors with 208 amino acid residues. FGF-6 is an important role in the regulation of cell proliferation, cell differentiation, angiogenesis and myogenesis. In physiological function, FGF6 is required for normal muscle regeneration.		
别称	fibroblast growth factor 6, HBGF-6, HST-2, FGF6		
蛋白编码	P10 <sup>7</sup> 67		
分子量	The protein has a calculated MW of 19.66 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).		
表达系统	Escherichia coli		
纯度	>98% as determined by SDS-PAGE.		
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED $_{50}$ for this effect is <0.1 ng/mL. The specific activity of recombinant human FGF-6 is > 1 x $10^7$ IU/mg.		
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.		
蛋白序列	MGTRANNTLLDSRGWGTLLSRSRAGLAGEIAGVNWESGYLVGIKRQRRLYCNVGIGFHLQVL PDGRISGTHEENPYSLLEISTVERGVVSLFGVRSALFVAMNSKGRLYATPSFQEECKFRETLLPNN YNAYESDLYQGTYIALSKYGRVKRGSKVSPIMTVTHFLPRI with polyhistidine tag at the Cterminus.		
产品形式	The protein was lyophilized from a 0.2 $\mu m$ filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.		
产品应用	Cell Culture		

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

储存 Initial Reconstitution:


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示



SDS-PAGE analysis of recombinant human FGF-6

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491