# **CHAMOT**

重组人成纤维细胞生长因子-11亚型2 /Recombinant FGF-11 isoform 2, Human, Animal-Free


CM101-5HP

CM101-20HP

CM101-100HP

CM101-500HP

CM101-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人成纤维细胞生长因子-11亚型2 /Recombinant FGF-11 isoform 2, Human, Animal-Free

| 产品编号 | CM101-5HP | CM101-20HP | CM101-100HP | CM101-500HP | CM101-1000HP |
|------|-----------|------------|-------------|-------------|--------------|
| 规格   | 5 μg      | 20 μg      | 100 μg      | 500 μg      | 1 mg         |

| 产品简介  |   |  |  |
|-------|---|--|--|
| 背景描述  | The Fibroblast Growth Factors (FGF) family has two group, secreted FGFs and intracellular FGFs (iFGFs). iFGFs are related to voltage-gated sodium (Nav) channels. Human FGF-11 isoform 2 is a 18 kDa cytokine with 166 amino acid residues. |  |  |
| 别称    | fibroblast growth factor 11 isoform 2, FHF-3, FHF3, FGF11   |  |  |
| 蛋白编码  | Q92914  |  |  |
| 分子量   | The protein has a calculated MW of 19.34 kDa. The protein migrates as 23 kDa under reducing condition (SDS-PAGE analysis).  |  |  |
| 表达系统  | Escherichia coli  |  |  |
| 纯度    | >98% as determined by SDS-PAGE.   |  |  |
| 生物活性  | Measure by its ability to induce 3T3 cells proliferation. The ED $_{50}$ for this effect is <1 ng/mL.   |  |  |
| 内毒素检测 | $<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.  |  |  |
| 蛋白序列  | MSLSPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIPVGLRVVTIQSAKLGHY MAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALYRQRRSGRAWYLGLDKEGQVMKGNR VKKTKAAAHFLPKLLEVAMYQEPSLHSVPEASPSSPPAP with polyhistidine tag at the Cterminus.                           |  |  |
| 产品形式  | The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.  |  |  |
| 产品应用  | Cell Culture  |  |  |

### 产品使用

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

储存

**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.


**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or


stabilizer.

运输

Blue Ice

### 实验数据展示





SDS-PAGE analysis of recombinant human FGF-

Human FGF-11 isoform 2, His Tag, E. coli induced 3T3 cell proliferation, with the ED50 at 0.84 ng/mL.

11 isoform 2

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491