# **CHAMOT**

重组人成纤维细胞生长因子-14 /Recombinant FGF-14, Human, Animal-Free

CM104-5HP

CM104-20HP

CM104-100HP

CM104-500HP

CM104-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



## **Product Datasheet**

# 重组人成纤维细胞生长因子-14 /Recombinant FGF-14, Human, Animal-Free

产品编号	CM104-5HP	CM104-20HP	CM104-100HP	CM104-500HP	CM104-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介			
背景描述	Fibroblast Growth Factors-14 (FGF-14) is a 27.7 kDa member of the fibroblast Growth Factors with 247 amino acid residues. FGF-14 is mainly expressed from brain, cervix. FGF-14 involved in nervous system development and function. May regulate voltage-gated sodium channels transport and function.		
别称	fibroblast growth factor 14, FHF-4, FHF4, SCA27, FGF14		
蛋白编码	Q92915		
分子量	The protein has a calculated MW of 28.28 kDa. The protein migrates as 33 kDa under reducing condition (SDS-PAGE analysis).		
表达系统	Escherichia coli		
纯度	>95% as determined by SDS-PAGE.		
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED $_{50}$ for this effect is <21 ng/mL.		
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.		
蛋白序列	AAAIASGLIRQKRQAREQHWDRPSASRRRSSPSKNRGLCNGNLVDIFSKVRIFGLKKRRLRRQ DPQLKGIVTRLYCRQGYYLQMHPDGALDGTKDDSTNSTLFNLIPVGLRVVAIQGVKTGLYIAM NGEGYLYPSELFTPECKFKESVFENYYVIYSSMLYRQQESGRAWFLGLNKEGQAMKGNRVKKT KPAAHFLPKPLEVAMYREPSLHDVGETVPKPGVTPSKSTSASAIMNGGKPVNKSKT with polyhistidine tag at the N-terminus.		
产品形式	The protein was lyophilized from a 0.2 $\mu m$ filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.		
产品应用	Cell Culture		

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

**Lyophilized Protein:** Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.


Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示





SDS-PAGE analysis of recombinant human FGF- Human FGF-14, His Tag, E. coli induced 3T3 cell proliferation, with

the ED50 at 20.35 ng/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491