CHAMOT

重组人成纤维细胞生长因子-23 /Recombinant FGF-23, Human, Animal-Free

CM112-5HP

CM112-20HP

CM112-100HP

CM112-500HP

CM112-1000HP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



≠品符介

背景描述

蛋白序列

Product Datasheet

重组人成纤维细胞生长因子-23 /Recombinant FGF-23, Human, Animal-Free

产品编号	CM112-5HP	CM112-20HP	CM112-100HP	CM112-500HP	CM112-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

/ HHIPJ/I	
	Fibroblast Growth Factors-23 (FGF-23) is a 28 kDa member of the fibroblast Growth
	Factors with 251 amino acid recidues EGE-23 is expressed from brain, benetic stellate

Factors with 251 amino acid residues. FGF-23 is expressed from brain, hepatic stellate cells, cone photoreceptor cells, early spermatids. FGF-23 involved phosphate metabolism and vitamin D metabolism. Negatively regulates osteoblast differentiation and matrix mineralization.

别称 fibroblast growth factor 23, FGFN, FGF23

蛋白编码 Q9GZV9

分子量 The protein has a calculated MW of 26.27 kDa. The protein migrates as 26 kDa under reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >98% as determined by SDS-PAGE.

生物活性 Measure by its ability to induce proliferation in BaF3 cells transfected with human FGFRIIIc. The ED50 for this effect is <0.3 µg/mL.

内毒素检测 <0.1 EU per 1 µg of the protein by the LAL method.

MYPNASPLLGSSWGGLIHLYTATARNSYHLQIHKNGHVDGAPHQTIYSALMIRSEDAGFVVIT GVMSRRYLCMDFRGNIFGSHYFDPENCRFQHQTLENGYDVYHSPQYHFLVSLGRAKRAFLPG MNPPPYSQFLSRRNEIPLIHFNTPIPRRHTRSAEDDSERDPLNVLKPRARMTPAPASCSQELPS AEDNSPMASDPLGVVRGGRVNTHAGGTGPEGCRPFAKFI with polyhistidine tag at the

C-terminus.

产品形式

The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.


Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示





SDS-PAGE analysis of recombinant human FGF-

Human FGF-23, His Tag, E. coli induced proliferation in BaF3 cells transfected with human FGFRIIIc, with the ED $_{50}$ at 0.26 μ g/mL.

23

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491