CHAMOT

重组人/小鼠/大鼠脑源性神经营养因子 /Recombinant BDNF, Human/Mouse/Rat, Animal-Free


CM135-5HP

CM135-20HP

CM135-100HP

CM135-500HP

CM135-1000HP





トス凹とつい

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

5 文献引用



Product Datasheet

重组人/小鼠/大鼠脑源性神经营养因子 /Recombinant BDNF, Human/Mouse/Rat, Animal-Free

产品编号	CM135-5HP	CM135-20HP	CM135-100HP	CM135-500HP	CM135-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介

产品应用

Cell Culture

背景描述	Brain-derived neurotrophic factor (BDNF) is a member of neurotrophin family that not only primarily expressed in hippocampus, amygdala, cerebral cortex, hypothalamus and cerebellum but also has been detected in blood platelets and in circulating plasma. BDNF is a 27.8 kDa protein containing 247 residues, which plays a critical role in regulating synaptic transmission and plasticity in various region of the CNS. Additionally, BNDF can acts as a modulator in the long-term potentiation of memory-related modifications in hippocampal synaptic transmission.		
别称	brain-derived neurotrophic factor, ANON2, BULN2		
蛋白编码	P23560(Human), P21237(Mouse), P23363(Rat)		
分子量	The protein has a calculated MW of 14.45 kDa. The protein migrates as 14 kDa under reducing condition (SDS-PAGE analysis).		
表达系统	Escherichia coli		
纯度	>98% as determined by SDS-PAGE.		
生物活性	Measure by its ability to induce proliferation in BaF3 cells transfected with TrkB. The ED $_{50}$ for this effect is <2 ng/mL.		
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.		
蛋白序列	MHSDPARRGELSVCDSISEWVTAADKKTAVDMSGGTVTVLEKVPVSKGQLKQYFYETKCNPMGYTKEGCRGIDKRHWNSQCRTTQSYVRALTMDSKKRIGWRFIRIDTSCVCTLTIKRGR with polyhistidine tag at the C-terminus.		
产品形式	The protein was lyophilized from a 0.2 μm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5. If you have any concerns or special requirements, please confirm with us.		

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



文献引用

• BDNF Alleviates Microglial Inhibition and Stereotypic Behaviors in a Mouse Model of Obsessive-Compulsive Disorder. Front Mol Neurosci. 2022 Jul 12;15:926572. (IF 6.261)(Human/Mouse/Rat BDNF)

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491