# **CHAMOT**

## 重组人胃内因子 /Recombinant GIF, Human, Animal-Free

CM144-5HP

CM144-20HP

CM144-100HP

CM144-500HP

CM144-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

### 重组人胃内因子 /Recombinant GIF, Human, Animal-Free

产品编号	CM144-5HP	CM144-20HP	CM144-100HP	CM144-500HP	CM144-1000HP
规格	5 μg	20 μg	100 µg	500 μg	1 mg

4	Е	7	<u>4-4</u>	•	Λ.
	П	ы	ľΞ	М	n

背景描述

蛋白序列

Human Gastric intrinsic factor (GIF) is an intrinsic factor of the eukaryotic cobalamin family. Gastric intrinsic factor is a 44 kDa protein containing 380 amino acid residues. It is secreted from the parietal cells in gastric mucosa. Gastric intrinsic factor plays a major role in absorption and transportation of cobalamin and Cbl. It could form the comprised with Vit.B12 that avoid the destroy of Vit.B12.

别称 gastric intrinsic factor, CBLIF, IF, IFMH, INF, TCN3

**蛋白编码** P27352

The protein has a calculated MW of 46.23 kDa. The protein migrates as 43 kDa under

reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >98% as determined by SDS-PAGE.

**生物活性** Testing in process

内毒素检测 <0.1 EU per 1 µg of the protein by the LAL method.

MAWFALYLLSLLWATAGTSTQTQSSCSVPSAQEPLVNGIQVLMENSVTSSAYPNPSILIAMNLA GAYNLKAQKLLTYQLMSSDNNDLTIGQLGLTIMALTSSCRDPGDKVSILQRQMENWAPSSPN AEASAFYGPSLAILALCQKNSEATLPIAVRFAKTLLANSSPFNVDTGAMATLALTCMYNKIPVGS EEGYRSLFGQVLKDIVEKISMKIKDNGIIGDIYSTGLAMQALSVTPEPSKKEWNCKKTTDMILNE IKQGKFHNPMSIAQILPSLKGKTYLDVPQVTCSPDHEVQPTLPSNPGPGPTSASNITVIYTINN

QLRGVELLFNETINVSVKSGSVLLVVLEEAQRKNPMFKFETTMTSWGLVVSSINNIAENVNHK
TYWQFLSGVTPLNEGVADYIPFNHEHITANFTQY with polyhistidine tag at the C-

terminus.

The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH

8.0. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

### 产品使用

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

**Lyophilized Protein:** Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491