

CHAMOT

**重组人半乳糖凝集素-3 /Recombinant Galectin-3,
Human, Animal-Free**

CM150-5HP

CM150-20HP

CM150-100HP

CM150-500HP

CM150-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人半乳糖凝集素-3 /Recombinant Galectin-3, Human, Animal-Free

产品编号	CM150-5HP	CM150-20HP	CM150-100HP	CM150-500HP	CM150-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Galectin-3 (Gal-3) is one of the lectin family members, which is the only chimera-type galectin, containing one carbohydrate recognition domain (CRD) connected to a long, flexible N-terminal domain. The C-terminal CRD is responsible for β-galactoside binding, and the N-terminal domain is essential for its multimerization, and interaction with other intracellular proteins. Galectin-3 is predominantly presented in the cytoplasm and expressed on the cell surface, and then often secreted into biological fluids, such as serum and urine. Numerous studies have indicated that galectin-3 plays a crucial role in cell proliferation, apoptosis, gene expression, immune surveillance, inflammation, fibrosis, and host defense.
别称	IgE-binding protein, MAC2, L-29, CPB-35
蛋白编码	P17931
分子量	The protein has a calculated MW of 27 kDa. The protein migrates as 32 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measured by its ability to chemoattract human THP1 cells using a concentration range of 0.5-1 µg/mL.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	ADNFSLHDALSGSGNPNPQGWPAGWGNQPAGAGGYPGASYPGAYPGQAPPGAYPGQAPP GAYPGAPGAYPGAPAGVYPGPPSGPGAYPSSGQPSATGAYPATGPYGAPAGPLIVPYNLPLPG GVVPRMLITILGTVKPNANRIALDFQRGNDVAFHFNPRFNENNRVIVCNTKLDNNWGREER QSVFPFESGKPFKIQVLVEPDHFKVAVNDAHLLQYNHRVKKLNEISKLGISGDIDLTSASYTMI with polyhistidine tag at the N-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human Galectin-3

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491