# **CHAMOT**

重组人半乳糖凝集素-7 /Recombinant Galectin-7, Human, Animal-Free

CM152-5HP

CM152-20HP

CM152-100HP

CM152-500HP

CM152-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人半乳糖凝集素-7 /Recombinant Galectin-7, Human, Animal-Free

产品编号	CM152-5HP	CM152-20HP	CM152-100HP	CM152-500HP	CM152-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

### 产品简介

背景描述	contains one carbohydrate recognition domain (CRD), responsible for $\beta$ -galactoside binding, and is biologically active as homodimers. Galectin-7 is constitutively presented in the gastrointestinal tract, stratified epithelia, skin, and fetal heart. It has been reported that galectin-7 serves as an effector in triggering apoptosis following activation of the p53 pathway. Furthermore, it serves important functions in numerous biological activities including cell adhesion, migration, proliferation, and differentiation.
别称	LGALS7B, LGALS7, PI7; GAL7, Gal-7, HKL-14
	activation of the p53 pathway. Furthermore, it serves important functions in numerous biological activities including cell adhesion, migration, proliferation, and differentiation.

蛋白编码	P47929
分子量	The protein has a calculated MW of 15.9 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).
<b>士</b> \上云 <i>は</i>	made autobita and the

表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
	Measure by its ability to chemoattract human THP1 cells. Chemotactic activity was

生物活性	observed at a concentration of $0.5 \mu g/mL$ .

< 0.1 EU per 1  $\mu g$  of the protein by the LAL method.

	蛋白序列	SNVPHKSSLPEGIRPGTVLRIRGLVPPNASRFHVNLLCGEEQGSDAALHFNPRLDTSEVVFNSK EQGSWGREERGPGVPFQRGQPFEVLIIASDDGFKAVVGDAQYHHFRHRLPLARVRLVEVGGD VQLDSVRIF with polyhistidine tag at the N-terminus.
--	------	---

产品形式	The protein was lyophilized from a 0.2 $\mu$ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品形式	

产品应用 Cell Culture

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491