# **CHAMOT**

重组人半乳糖凝集素-13 /Recombinant Galectin-13, Human, Animal-Free

CM157-5HP

CM157-20HP

CM157-100HP

CM157-500HP

CM157-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人半乳糖凝集素-13 /Recombinant Galectin-13, Human, Animal-Free

产品编号	CM157-5HP	CM157-20HP	CM157-100HP	CM157-500HP	CM157-1000HP
规格	5 μg	20 μg	100 µg	500 μg	1 mg

### 产品简介

背景描述

Galectin-13 (Gal-13) is a lectin family member and is one of the prototype galectins. It contains one carbohydrate recognition domain (CRD), responsible for  $\beta$ -galactoside binding, and is biologically active as homodimers. Galectin-13 is expressed in the placenta, contributing to T cell apoptosis, immune regulation, and immune tolerance, which is indispensable for fetal development. For example, galectin-13 organizes crystal-like aggregates in the decidua, consequently attracting and killing maternal immune cells. Furthermore, galectin-13 is essential for driving neutrophil polarization in the placental-growth-permissive phenotype.

	neutrophil polarization in the placental-growth-permissive phenotype.
别称	LGALS13, GAL13, PLAC8, PP13
蛋白编码	Q9UHV8
分子量	The protein has a calculated MW of 16.9 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Testing in process
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.
蛋白序列	SSLPVPYKLPVSLSVGSCVIIKGTPIHSFINDPQLQVDFYTDMDEDSDIAFRFRVHFGNHVVMN RREFGIWMLEETTDYVPFEDGKQFELCIYVHYNEYEIKVNGIRIYGFVHRIPPSFVKMVQVSRDI SLTSVCVCN with polyhistidine tag at the N-terminus.

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

7.4. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

产品形式

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

**Lyophilized Protein:** Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

### 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491